



The Riemann Hypothesis

Frank Vega

EasyChair preprints are intended for rapid dissemination of research results and are integrated with the rest of EasyChair.

August 25, 2020

The Riemann hypothesis

Frank Vega

Abstract. In mathematics, the Riemann hypothesis is a conjecture that the Riemann zeta function has its zeros only at the negative even integers and complex numbers with real part $\frac{1}{2}$. Many consider it to be the most important unsolved problem in pure mathematics. The Robin's inequality consists in $\sigma(n) < e^\gamma \times n \times \ln \ln n$ where $\sigma(n)$ is the divisor function and $\gamma \approx 0.57721$ is the Euler-Mascheroni constant. The Robin's inequality is true for every natural number $n > 5040$ if and only if the Riemann hypothesis is true. We demonstrate the Robin's inequality is true for every natural number $n > 5040$. Consequently, we show the Riemann hypothesis is true.

1 Introduction

In mathematics, the Riemann hypothesis is a conjecture that the Riemann zeta function has its zeros only at the negative even integers and complex numbers with real part $\frac{1}{2}$. Many consider it to be the most important unsolved problem in pure mathematics [5]. It is of great interest in number theory because it implies results about the distribution of prime numbers [5]. It was proposed by Bernhard Riemann (1859), after whom it is named [5]. It is one of the seven Millennium Prize Problems selected by the Clay Mathematics Institute to carry a US 1,000,000 prize for the first correct solution [5]. The divisor function $\sigma(n)$ for a natural number n is defined as the sum of the powers of the divisors of n ,

$$\sigma(n) = \sum_{k|n} k$$

where $k | n$ means that the natural number k divides n [6]. In 1915, Ramanujan proved that under the assumption of the Riemann hypothesis, the inequality,

$$\sigma(n) < e^\gamma \times n \times \ln \ln n$$

holds for all sufficiently large n , where $\gamma \approx 0.57721$ is the Euler-Mascheroni constant [3]. The largest known value that violates the inequality is $n = 5040$. In 1984, Guy Robin proved that the inequality is true for all $n > 5040$ if and only if the Riemann hypothesis is true [3]. Using this inequality, we show that the Riemann hypothesis is true.

2010 Mathematics Subject Classification: Primary 11M26.

Keywords: number theory, inequality, divisor, prime.

2 Results

Theorem 2.1 Given a natural number $n = p_1^{a_1} \times p_2^{a_2} \times \dots \times p_m^{a_m}$ such that p_1, p_2, \dots, p_m are prime numbers, then we obtain the following inequality

$$\frac{\sigma(n)}{n} < \prod_{i=1}^m \frac{p_i}{p_i - 1}.$$

Proof For a natural number $n = p_1^{a_1} \times p_2^{a_2} \times \dots \times p_m^{a_m}$ such that p_1, p_2, \dots, p_m are prime numbers, then we obtain the following formula

$$(2.1) \quad \sigma(n) = \prod_{i=1}^m \frac{p_i^{a_i+1} - 1}{p_i - 1}$$

from the Ramanujan's notebooks [1]. In this way, we have that

$$\frac{\sigma(n)}{n} = \prod_{i=1}^m \frac{p_i^{a_i+1} - 1}{p_i^{a_i} \times (p_i - 1)}.$$

However, for any prime power $p_i^{a_i}$, we have that

$$\frac{p_i^{a_i+1} - 1}{p_i^{a_i} \times (p_i - 1)} < \frac{p_i^{a_i+1}}{p_i^{a_i} \times (p_i - 1)} = \frac{p_i}{p_i - 1}.$$

Consequently, we obtain that

$$\frac{\sigma(n)}{n} < \prod_{i=1}^m \frac{p_i}{p_i - 1}.$$

■

Theorem 2.2 Given some prime numbers p_1, p_2, \dots, p_m , then we obtain the following inequality,

$$\prod_{i=1}^m \frac{p_i}{p_i - 1} < \frac{\pi^2}{6} \times \prod_{i=1}^m \frac{p_i + 1}{p_i}.$$

Proof Given a prime number p_i , we obtain that

$$\frac{p_i}{p_i - 1} = \frac{p_i^2}{p_i^2 - p_i}$$

and that would be equivalent to

$$\frac{p_i^2}{p_i^2 - p_i} = \frac{p_i^2}{p_i^2 - 1 - (p_i - 1)}$$

and that is the same as

$$\frac{p_i^2}{p_i^2 - 1 - (p_i - 1)} = \frac{p_i^2}{(p_i - 1) \times \left(\frac{p_i^2 - 1}{p_i - 1} - 1\right)}$$

which is equal to

$$\frac{p_i^2}{(p_i - 1) \times \left(\frac{p_i^2 - 1}{(p_i - 1)} - 1\right)} = \frac{p_i^2}{(p_i - 1) \times \frac{p_i^2 - 1}{(p_i - 1)} \times \left(1 - \frac{(p_i - 1)}{p_i^2 - 1}\right)}$$

that is equivalent to

$$\frac{p_i^2}{(p_i - 1) \times \frac{p_i^2 - 1}{(p_i - 1)} \times \left(1 - \frac{(p_i - 1)}{p_i^2 - 1}\right)} = \frac{p_i^2}{p_i^2 - 1} \times \frac{1}{1 - \frac{(p_i - 1)}{p_i^2 - 1}}$$

which is the same as

$$\frac{p_i^2}{p_i^2 - 1} \times \frac{1}{1 - \frac{(p_i - 1)}{p_i^2 - 1}} = \frac{1}{1 - p_i^{-2}} \times \frac{1}{1 - \frac{1}{(p_i + 1)}}$$

and finally

$$\frac{1}{(1 - p_i^{-2})} \times \frac{1}{1 - \frac{1}{(p_i + 1)}} = \frac{1}{(1 - p_i^{-2})} \times \frac{p_i + 1}{p_i}.$$

In this way, we have that

$$\prod_{i=1}^m \frac{p_i}{p_i - 1} = \prod_{i=1}^m \frac{1}{1 - p_i^{-2}} \times \prod_{i=1}^m \frac{p_i + 1}{p_i}.$$

However, we know that

$$\prod_{i=1}^m \frac{1}{1 - p_i^{-2}} < \prod_{j=1}^{\infty} \frac{1}{1 - p_j^{-2}}$$

where p_j is the j^{th} prime number and we have that

$$\prod_{j=1}^{\infty} \frac{1}{1 - p_j^{-2}} = \frac{\pi^2}{6}$$

as a consequence of the result in the Basel problem [6]. Consequently, we obtain that

$$\prod_{i=1}^m \frac{p_i}{p_i - 1} < \frac{\pi^2}{6} \times \prod_{i=1}^m \frac{p_i + 1}{p_i}.$$

■

Definition 2.3 We recall that an integer n is said to be squarefree if for every prime divisor p of n we have $p^2 \nmid n$, where $p^2 \nmid n$ means that p^2 does not divide n [2].

Theorem 2.4 Given a squarefree number $n = q_1 \times \dots \times q_m$ such that the greatest prime divisor of n is greater than 3, then we obtain the following inequality

$$\sigma(n) < \frac{6}{\pi^2} \times e^{\gamma} \times n \times \ln \ln n.$$

Proof This proof is very similar with the demonstration in Theorem 1.1 from the article reference [2]. By induction with respect to $\omega(n)$, that is the number of distinct prime factors of n [2]. Put $\omega(n) = m$ [2]. We need to prove the assertion for those integers with $m = 1$. From the formula (2.1), we obtain that

$$\sigma(n) = (p_1 + 1) \times (p_2 + 1) \times \dots \times (p_m + 1)$$

when $n = p_1 \times p_2 \times \dots \times p_m$. In this way, for any prime number $p_i \geq 5$, then we need to prove

$$(2.2) \quad \left(1 + \frac{1}{p_i}\right) < \frac{6}{\pi^2} \times e^\gamma \times \ln \ln(p_i).$$

For $p_i = 5$, we have that

$$\left(1 + \frac{1}{5}\right) < \frac{6}{\pi^2} \times e^\gamma \times \ln \ln(5)$$

is actually true. For another prime number $p_i > 5$, we have that

$$\left(1 + \frac{1}{p_i}\right) < \left(1 + \frac{1}{5}\right)$$

and

$$\frac{6}{\pi^2} \times e^\gamma \times \ln \ln(5) < \frac{6}{\pi^2} \times e^\gamma \times \ln \ln(p_i)$$

which clearly implies that the inequality (2.2) is true for every prime number $p_i \geq 5$. Now, suppose it is true for $m - 1$, with $m \geq 1$ and let us consider the assertion for those squarefree n with $\omega(n) = m$ [2]. So let $n = q_1 \times \dots \times q_m$ be a squarefree number and assume that $q_1 < \dots < q_m$ for $q_m \geq 5$.

Case1 : $q_m \geq \ln(q_1 \times \dots \times q_{m-1} \times q_m) = \ln n$ for $q_m \geq 5$.

By the induction hypothesis we have

$$(q_1 + 1) \times \dots \times (q_{m-1} + 1) < \frac{6}{\pi^2} \times e^\gamma \times q_1 \times \dots \times q_{m-1} \times \ln \ln(q_1 \times \dots \times q_{m-1}),$$

and hence

$$(q_1 + 1) \times \dots \times (q_{m-1} + 1) \times (q_m + 1) < \frac{6}{\pi^2} \times e^\gamma \times q_1 \times \dots \times q_{m-1} \times (q_m + 1) \times \ln \ln(q_1 \times \dots \times q_{m-1})$$

when we multiply the both sides of the inequality by $(q_m + 1)$. We want to show that

$$\frac{6}{\pi^2} \times e^\gamma \times q_1 \times \dots \times q_{m-1} \times (q_m + 1) \times \ln \ln(q_1 \times \dots \times q_{m-1}) \leq \frac{6}{\pi^2} \times e^\gamma \times q_1 \times \dots \times q_{m-1} \times q_m \times \ln \ln(q_1 \times \dots \times q_{m-1} \times q_m) = \frac{6}{\pi^2} \times e^\gamma \times n \times \ln \ln n.$$

Indeed the previous inequality is equivalent with

$$q_m \times \ln \ln(q_1 \times \dots \times q_{m-1} \times q_m) \geq (q_m + 1) \times \ln \ln(q_1 \times \dots \times q_{m-1}),$$

or alternatively

$$\frac{q_m \times (\ln \ln(q_1 \times \dots \times q_{m-1} \times q_m) - \ln \ln(q_1 \times \dots \times q_{m-1}))}{\ln q_m} \geq$$

$$\frac{\ln \ln(q_1 \times \dots \times q_{m-1})}{\ln q_m}.$$

From the reference [2], we have that if $0 < a < b$, then

$$(2.3) \quad \frac{\ln b - \ln a}{b - a} = \frac{1}{(b - a)} \int_a^b \frac{dt}{t} > \frac{1}{b}.$$

We can apply the inequality (2.3) to the previous one just using $b = \ln(q_1 \times \dots \times q_{m-1} \times q_m)$ and $a = \ln(q_1 \times \dots \times q_{m-1})$. Certainly, we have that

$$\begin{aligned} \ln(q_1 \times \dots \times q_{m-1} \times q_m) - \ln(q_1 \times \dots \times q_{m-1}) &= \\ \ln \frac{q_1 \times \dots \times q_{m-1} \times q_m}{q_1 \times \dots \times q_{m-1}} &= \ln q_m. \end{aligned}$$

In this way, we obtain that

$$\frac{q_m \times (\ln \ln(q_1 \times \dots \times q_{m-1} \times q_m) - \ln \ln(q_1 \times \dots \times q_{m-1}))}{\ln q_m} > \frac{q_m}{\ln(q_1 \times \dots \times q_m)}.$$

Using this result we infer that the original inequality is certainly satisfied if the next inequality is satisfied

$$\frac{q_m}{\ln(q_1 \times \dots \times q_m)} \geq \frac{\ln \ln(q_1 \times \dots \times q_{m-1})}{\ln q_m}$$

which is trivially true for $q_m \geq \ln(q_1 \times \dots \times q_{m-1} \times q_m)$ [2].

Case2 : $q_m < \ln(q_1 \times \dots \times q_{m-1} \times q_m) = \ln n$ for $q_m \geq 5$.

Since, we need to prove that

$$\frac{\sigma(n)}{n} < \frac{6}{\pi^2} \times e^\gamma \times \ln \ln n$$

that would be the same as

$$(2.4) \quad \prod_{i=1}^m \frac{p_i}{p_i - 1} < \frac{6}{\pi^2} \times e^\gamma \times \ln \ln n$$

according to Theorem 2.1. However, we know that

$$\prod_{p \leq x} \frac{p}{p-1} < \frac{\epsilon}{\ln x} \times \left(1 + \frac{1}{2 \times \ln^2 x}\right)$$

for $x \geq 1$, where $\epsilon < 0.561459483566886$ [4]. That would be the same as

$$\prod_{p \leq x} \frac{p}{p-1} < \frac{\epsilon}{\ln^3 x} \times \left(\ln^2 x + \frac{1}{2}\right)$$

and therefore, we would only need to check for the inequality (2.4) that

$$\frac{\epsilon}{\ln^3 q_m} \times \left(\ln^2 q_m + \frac{1}{2}\right) < \frac{6}{\pi^2} \times e^\gamma \times \ln \ln n$$

that is

$$\epsilon \times \left(\ln^2 q_m + \frac{1}{2}\right) < \frac{6}{\pi^2} \times e^\gamma \times \ln \ln n \times \ln^3 q_m$$

because of

$$\prod_{i=1}^m \frac{p_i}{p_i - 1} \leq \prod_{p \leq p_m} \frac{p}{p - 1}.$$

Under our assumption, we have that

$$\ln \ln n > \ln q_m$$

and therefore

$$\frac{6}{\pi^2} \times e^\gamma \times \ln^4 q_m < \frac{6}{\pi^2} \times e^\gamma \times \ln \ln n \times \ln^3 q_m$$

and thus, we only need to prove that

$$\epsilon \times \left(\ln^2 q_m + \frac{1}{2} \right) < \frac{6}{\pi^2} \times e^\gamma \times \ln^4 q_m.$$

We have that $\frac{6}{\pi^2} \times e^\gamma > 1.08276219326$ and thus

$$\frac{6}{\pi^2} \times e^\gamma \times \frac{1}{\epsilon} > \frac{1.08276219326}{0.561459483566886} > 1.927.$$

. Hence, we only need to prove that

$$\ln^2 q_m + \frac{1}{2} < 1.927 \times \ln^4 q_m.$$

In addition, we can note the function $f(x) = 1.927 \times \ln^4 x - \ln^2 x - \frac{1}{2}$ is strictly increasing. Under the notion of

$$1.927 \times \ln^4 q_m - \ln^2 q_m - \frac{1}{2} > 1.927 \times \ln^4 5 - \ln^2 5 - \frac{1}{2}$$

for $q_m \geq 5$, we only need to prove that

$$1.927 \times \ln^4 5 - \ln^2 5 - \frac{1}{2} > 0$$

which is true and this finally implies the Theorem is indeed satisfied. ■

Definition 2.5 We recall that an integer n is said to be squarefull if for every prime divisor p of n we have $p^2 \mid n$ [2].

Theorem 2.6 Given a natural number $n = p_1^{a_1} \times p_2^{a_2} \times \dots \times p_m^{a_m} > 5040$ such that p_1, p_2, \dots, p_m are prime numbers and the greatest prime divisor of n is lesser than or equal to 3, then the Robin's inequality is true for n .

Proof Given a natural number $n = p_1^{a_1} \times p_2^{a_2} \times \dots \times p_m^{a_m} > 5040$ such that p_1, p_2, \dots, p_m are prime numbers, we need to prove that

$$\frac{\sigma(n)}{n} < e^\gamma \times \ln \ln n$$

that would be the same as

$$(2.5) \quad \prod_{i=1}^m \frac{p_i}{p_i - 1} < e^\gamma \times \ln \ln n$$

according to Theorem 2.1. Since the Robin's inequality has been proved for every squarefull $n > 5040$ [2], then this would be true for $n = p_1^{a_1} > 5040$ and

$p_1 \leq 3$. Therefore, we only need to prove this for $n = 2^{a_1} \times 3^{a_2} > 5040$. In this way, we have that

$$\frac{2 \times 3}{1 \times 2} = 3 < e^\gamma \times \ln \ln(5040) \approx 3.81.$$

However, we know for $n > 5040$, we have that

$$e^\gamma \times \ln \ln(5040) < e^\gamma \times \ln \ln n$$

and thus, the proof is completed. \blacksquare

Theorem 2.7 *The Robin's inequality is true for every natural number $n > 5040$.*

Proof Given a natural number $n = p_1^{a_1} \times p_2^{a_2} \times \dots \times p_m^{a_m} > 5040$ such that p_1, p_2, \dots, p_m are prime numbers, then we will prove the Robin's inequality is true for n . We know this is true when the greatest prime divisor of n is lesser than or equal to 3 according to the Theorem 2.6. Consequently, we have to demonstrate when the greatest prime divisor of n is greater than 3. We need to prove that

$$\sigma(n) < e^\gamma \times n \times \ln \ln n$$

which is the same as

$$\frac{\sigma(n)}{n} < e^\gamma \times \ln \ln n.$$

However, we know that would be true when (2.5) is true according to Theorem 2.1. In addition, the inequality (2.5) would be true when

$$\frac{\pi^2}{6} \times \prod_{i=1}^m \frac{p_i + 1}{p_i} < e^\gamma \times \ln \ln n$$

according to Theorem 2.2. Using the properties of the formula (2.1), we obtain that will be equivalent to

$$\frac{\pi^2}{6} \times \frac{\sigma(n')}{n'} < e^\gamma \times \ln \ln n$$

where $n' = q_1 \times \dots \times q_m$ is a squarefree number. However, that would be equal to

$$\sigma(n') < \frac{6}{\pi^2} \times e^\gamma \times n' \times \ln \ln n.$$

In addition, if the greatest prime divisor of n is greater than 3, then the greatest prime divisor of n' is greater than 3 as well. Consequently, we have that

$$\frac{6}{\pi^2} \times e^\gamma \times n' \times \ln \ln n' < \frac{6}{\pi^2} \times e^\gamma \times n' \times \ln \ln n$$

and therefore, we only need to prove that

$$\sigma(n') < \frac{6}{\pi^2} \times e^\gamma \times n' \times \ln \ln n'$$

which is true according to the Theorem 2.4 when the greatest prime divisor of n' is greater than 3. To sum up, we have finally proved the Robin's inequality is true for every natural number $n > 5040$. \blacksquare

Theorem 2.8 *The Riemann hypothesis is true.*

Proof If the Robin's inequality is true for every natural number $n > 5040$, then the Riemann hypothesis is true [3]. Hence, the Riemann hypothesis is true due to Theorem 2.7. ■

3 Conclusions

The practical uses of the Riemann hypothesis include many propositions known true under the Riemann hypothesis, and some that can be shown equivalent to the Riemann hypothesis [5]. Certainly, the Riemann hypothesis is close related to various mathematical topics such as the distribution of prime numbers, the growth of arithmetic functions, the Lindelöf hypothesis, the large prime gap conjecture, etc [5]. In this way, a proof of the Riemann hypothesis could spur considerable advances in many mathematical areas, such as the number theory and pure mathematics [5].

References

- [1] Bruce C Berndt. *Ramanujan's notebooks*. Springer Science & Business Media, 2012.
- [2] YoungJu Choie, Nicolas Lichiardopol, Pieter Moree, and Patrick Solé. On Robin's criterion for the Riemann hypothesis. *Journal de théorie des nombres de Bordeaux*, 19(2):357–372, 2007.
- [3] Jeffrey C. Lagarias. An elementary problem equivalent to the riemann hypothesis. *The American Mathematical Monthly*, 109(6):534–543, 2002.
- [4] J Barkley Rosser and Lowell Schoenfeld. Approximate formulas for some functions of prime numbers. *Illinois Journal of Mathematics*, 6(1):64–94, 1962.
- [5] Peter Sarnak. Problems of the millennium: The riemann hypothesis (2004), April 2005. In Clay Mathematics Institute at http://www.claymath.org/library/annual_report/ar2004/04report_prizeproblem.pdf. Retrieved 24 August, 2020.
- [6] David G. Wells. *Prime Numbers, The Most Mysterious Figures in Math*. John Wiley & Sons, Inc., 2005.