



## On the Nontrivial Zeros of the Riemann Zeta Function

---

Frank Vega

EasyChair preprints are intended for rapid dissemination of research results and are integrated with the rest of EasyChair.

October 28, 2022

# On the nontrivial zeros of the Riemann zeta function

Frank Vega<sup>1\*</sup>

<sup>1\*</sup>Software Department, CopSonic, 1471 Route de Saint-Nauphary, Montauban, 82000, Tarn-et-Garonne, France.

Corresponding author(s). E-mail(s): [vega.frank@gmail.com](mailto:vega.frank@gmail.com);

## Abstract

The Riemann hypothesis is a conjecture that the Riemann zeta function has its zeros only at the negative even integers and complex numbers with real part  $\frac{1}{2}$ . It is considered by many to be the most important unsolved problem in pure mathematics. There are several statements equivalent to the famous Riemann hypothesis. In 2011, Solé and Planat stated that, the Riemann hypothesis is true if and only if the inequality  $\zeta(2) \cdot \prod_{p \leq x} (1 + \frac{1}{p}) > e^\gamma \cdot \log \theta(x)$  holds for all  $x \geq 5$ , where  $\theta(x)$  is the first Chebyshev function,  $\gamma \approx 0.57721$  is the Euler-Mascheroni constant,  $\zeta(x)$  is the Riemann zeta function and  $\log$  is the natural logarithm. In this note, using Solé and Planat criterion, we prove that, when the Riemann hypothesis is false, then there are infinitely many natural numbers  $x$  for which  $\frac{\log x}{\sqrt{x}} + O(\frac{1}{\sqrt{x}}) + 2 \cdot \log x + c \cdot \log x \leq 2.062$  could be satisfied for some  $c > 0$ . Since the inequality  $\frac{\log x}{\sqrt{x}} + O(\frac{1}{\sqrt{x}}) + 2 \cdot \log x + c \cdot \log x \leq 2.062$  does not hold for some  $c > 0$  and large enough  $x$ , then the Riemann hypothesis is true by principle of non-contradiction.

**Keywords:** Riemann hypothesis, Riemann zeta function, Chebyshev function

**MSC Classification:** 11M26 , 11A25

## 1 Introduction

The Riemann hypothesis is considered by many to be the most important unsolved problem in pure mathematics. It was proposed by Bernhard Riemann (1859). The Riemann hypothesis belongs to the Hilbert's eighth problem

## A Millennium Prize Problem

### 2 The Riemann hypothesis

on David Hilbert's list of twenty-three unsolved problems. Leonhard Euler discovered a particular value of the Riemann zeta function (1734).

**Proposition 1** *It is known that*[1, (1) p. 1070]:

$$\zeta(2) = \prod_{i=1}^{\infty} \frac{p_i^2}{p_i^2 - 1} = \frac{\pi^2}{6}.$$

**Proposition 2** *For*  $x > 24317$ :

$$\sum_{p>x} \log \left( \frac{p^2}{p^2 - 1} \right) = \frac{1}{x \cdot \log x} + O \left( \frac{1}{x \cdot \log^2 x} \right)$$

where  $\log$  is the natural logarithm [2].

In mathematics, the first Chebyshev function  $\theta(x)$  is given by

$$\theta(x) = \sum_{p \leq x} \log p$$

with the sum extending over all prime numbers  $p$  that are less than or equal to  $x$ . For  $x \geq 2$ , we say that  $\text{Dedekind}(x)$  holds provided that

$$\zeta(2) \cdot \prod_{p \leq x} \left( 1 + \frac{1}{p} \right) > e^\gamma \cdot \log \theta(x)$$

where  $\gamma \approx 0.57721$  is the Euler-Mascheroni constant. Next, we have Solé and Planat Theorem:

**Proposition 3**  $\text{Dedekind}(x)$  holds for all natural numbers  $x \geq 5$  if and only if the Riemann hypothesis is true [3, Theorem 4.2 p. 5].

This is the main insight.

**Lemma 1** *If the Riemann hypothesis is false, then there are infinitely many natural numbers*  $x \geq 5$  *for which*  $\text{Dedekind}(x)$  *fails (i.e. Dedekind}(x) does not hold).*

Srinivasa Ramanujan studied the function  $S_1(x) = \sum_{\rho} \frac{x^{\rho-1}}{\rho \cdot (1-\rho)}$  where  $\rho$  runs over the nontrivial zeros of the Riemann  $\zeta$  function [4, Section 65]. In number theory, the second Chebyshev function  $\psi(x)$  is given by

$$\psi(x) = \sum_{p^k \leq x} \log p$$

with the sum extending over all prime powers  $p^k$  that are less than or equal to  $x$ . Thus, we have two Nicolas Theorems:

**Proposition 4** For some  $c > 0$ :

$$\begin{aligned} S_1(x) &= - \int_1^x \frac{\psi(t)}{t^2} dt + \log x - 1 - \gamma + \frac{\log(2 \cdot \pi)}{x} - \sum_{k=1}^{\infty} \frac{1}{2 \cdot k \cdot (2 \cdot k + 1) \cdot x^{2 \cdot k + 1}} \\ &= \frac{c}{\sqrt{x}} \end{aligned}$$

for every  $x > 1$  [2].

**Proposition 5** [5, (2.16)]. For all  $x \geq 10^9$ :

$$\sum_{p \leq x} \log \left( 1 - \frac{1}{p} \right) + \gamma + \log \log \theta(x) + \frac{2}{\sqrt{x} \cdot \log x} + \frac{S_1(x)}{\log x} \leq \frac{2.062}{\sqrt{x} \cdot \log^2 x}$$

This is our main theorem.

**Theorem 1** If the Riemann hypothesis is false, then there are infinitely many natural numbers  $x$  for which

$$\frac{\log x}{\sqrt{x}} + O \left( \frac{1}{\sqrt{x}} \right) + 2 \cdot \log x + c \cdot \log x \leq 2.062$$

could be satisfied for some  $c > 0$ .

The following is a key Corollary.

**Corollary 1** The Riemann hypothesis is true.

## 2 Proof of the Lemma 1

*Proof* According to Proposition 3, the Riemann hypothesis is false, if there exists some natural number  $x_0 \geq 5$  such that  $g(x_0) > 1$  or equivalent  $\log g(x_0) > 0$ :

$$g(x) = \frac{e^\gamma}{\zeta(2)} \cdot \log \theta(x) \cdot \prod_{p \leq x} \left( 1 + \frac{1}{p} \right)^{-1}.$$

We know the bound [3, Theorem 4.2 p. 5]:

$$\log g(x) \geq \log f(x) - \frac{2}{x}$$

where  $f$  was introduced in the Nicolas paper [6, Theorem 3 p. 376]:

$$f(x) = e^\gamma \cdot \log \theta(x) \cdot \prod_{p \leq x} \left( 1 - \frac{1}{p} \right).$$

4 *The Riemann hypothesis*

When the Riemann hypothesis is false, then there exists a real number  $b < \frac{1}{2}$  for which there are infinitely many natural numbers  $x$  such that  $\log f(x) = \Omega_+(x^{-b})$  [6, Theorem 3 (c) p. 376]. According to the Hardy and Littlewood definition, this would mean that

$$\exists k > 0, \forall y_0 \in \mathbb{N}, \exists y \in \mathbb{N} (y > y_0): \log f(y) \geq k \cdot y^{-b}.$$

That inequality is equivalent to  $\log f(y) \geq (k \cdot y^{-b} \cdot \sqrt{y}) \cdot \frac{1}{\sqrt{y}}$ , but we note that

$$\lim_{y \rightarrow \infty} (k \cdot y^{-b} \cdot \sqrt{y}) = \infty$$

for every possible positive value of  $k$  when  $b < \frac{1}{2}$ . In this way, this implies that

$$\forall y_0 \in \mathbb{N}, \exists y \in \mathbb{N} (y > y_0): \log f(y) \geq \frac{1}{\sqrt{y}}.$$

Hence, if the Riemann hypothesis is false, then there are infinitely many natural numbers  $x$  such that  $\log f(x) \geq \frac{1}{\sqrt{x}}$ . Since  $\frac{2}{x} = o(\frac{1}{\sqrt{x}})$ , then it would be infinitely many natural numbers  $x_0$  such that  $\log g(x_0) > 0$ .  $\square$

### 3 Proof of the Theorem 1

By Lemma 1, there are infinitely many natural numbers  $x \geq 5$  for which

$$\gamma + \log \log \theta(x) \geq \sum_{p \leq x} \log \left( 1 + \frac{1}{p} \right) + \log(\zeta(2))$$

could be satisfied. This implies that there are infinitely many natural numbers  $x \geq 10^9$  for which

$$\sum_{p \leq x} \log \left( 1 - \frac{1}{p} \right) + \sum_{p \leq x} \log \left( 1 + \frac{1}{p} \right) + \log(\zeta(2)) + \frac{2}{\sqrt{x} \cdot \log x} + \frac{S_1(x)}{\log x} \leq \frac{2.062}{\sqrt{x} \cdot \log^2 x}$$

could be also satisfied by Proposition 5. We can write

$$\left( 1 - \frac{1}{p} \right) \cdot \left( 1 + \frac{1}{p} \right) = \left( 1 - \frac{1}{p^2} \right).$$

for every prime [3, p. 3]. Consequently, we obtain that

$$\sum_{p \leq x} \log \left( 1 - \frac{1}{p^2} \right) + \log(\zeta(2)) + \frac{2}{\sqrt{x} \cdot \log x} + \frac{S_1(x)}{\log x} \leq \frac{2.062}{\sqrt{x} \cdot \log^2 x}$$

by properties of logarithms. By Proposition 1, we know that

$$\sum_{p > x} \log \left( \frac{p^2}{p^2 - 1} \right) + \frac{2}{\sqrt{x} \cdot \log x} + \frac{S_1(x)}{\log x} \leq \frac{2.062}{\sqrt{x} \cdot \log^2 x}.$$

Using the Propositions 2 and 4, we can see that

$$\frac{1}{x \cdot \log x} + O\left(\frac{1}{x \cdot \log^2 x}\right) + \frac{2}{\sqrt{x} \cdot \log x} + \frac{c}{\sqrt{x} \cdot \log x} \leq \frac{2.062}{\sqrt{x} \cdot \log^2 x}.$$

Let's multiply both sides by  $\sqrt{x} \cdot \log^2 x$  to assume that

$$\frac{\log x}{\sqrt{x}} + O\left(\frac{1}{\sqrt{x}}\right) + 2 \cdot \log x + c \cdot \log x \leq 2.062$$

could be satisfied for some  $c > 0$ , when the Riemann hypothesis is false. Note that, there are not anomalies of signs after of multiplying by  $\sqrt{x} \cdot \log^2 x$  since for  $x \geq 10^9$  we obtain that  $\log x > 20.7$ .

## 4 Proof of Corollary 1

*Proof* By Theorem 1, we proved that there exists some  $c'$  such that there are infinitely many natural numbers  $x$  for which

$$\frac{\log x}{\sqrt{x}} + \frac{c'}{\sqrt{x}} + 2 \cdot \log x + c \cdot \log x \leq 2.062$$

could be satisfied for some  $c > 0$ , when the Riemann hypothesis is false. However, no matter how small could be the value of  $c'$ , we can always assure that

$$\frac{\log x}{\sqrt{x}} + \frac{c'}{\sqrt{x}} + 2 \cdot \log x + c \cdot \log x \leq 2.062$$

does not hold for some  $c > 0$  and large enough  $x$ . Note that, the Theorem 1 was stated by the domain of the natural numbers, so that is the same to say that there exists some  $y > 0$  such that for all natural numbers  $x > y$ , we can confirm that the inequality does not hold. In conclusion, the Riemann hypothesis is true by principle of non-contradiction. □

## 5 Conclusions

Practical uses of the Riemann hypothesis include many propositions that are known to be true under the Riemann hypothesis and some that can be shown to be equivalent to the Riemann hypothesis. Indeed, the Riemann hypothesis is closely related to various mathematical topics such as the distribution of primes, the growth of arithmetic functions, the Lindelöf hypothesis, the Large Prime Gap Conjecture, etc. Certainly, a proof of the Riemann hypothesis could spur considerable advances in many mathematical areas, such as number theory and pure mathematics in general.

**Acknowledgments.** The author wishes to thank his mother, maternal brother, maternal aunt, and friends Liuva, Yary, Sonia, and Arelis for their support.

## References

- [1] R. Ayoub, Euler and the zeta function. *The American Mathematical Monthly* **81**(10), 1067–1086 (1974). <https://doi.org/10.2307/2319041>
- [2] J.L. Nicolas, The sum of divisors function and the Riemann hypothesis. *The Ramanujan Journal* **58**, 1113–1157 (2022). <https://doi.org/10.1007/s11139-021-00491-y>
- [3] P. Solé, M. Planat, Extreme values of the Dedekind  $\psi$  function. *Journal of Combinatorics and Number Theory* **3**(1), 33–38 (2011)
- [4] J.L. Nicolas, G. Robin, Highly Composite Numbers by Srinivasa Ramanujan. *The Ramanujan Journal* **1**(2), 119–153 (1997). <https://doi.org/10.1023/A:1009764017495>
- [5] J.L. Nicolas, Small values of the Euler function and the Riemann hypothesis. *Acta Arithmetica* **155**, 311–321 (2012). <https://doi.org/10.4064/aa155-3-7>
- [6] J.L. Nicolas, Petites valeurs de la fonction d’Euler. *Journal of number theory* **17**(3), 375–388 (1983). [https://doi.org/10.1016/0022-314X\(83\)90055-0](https://doi.org/10.1016/0022-314X(83)90055-0)